

EXTRACTION POST-OP INSTRUCTIONS

REMEMBER THAT YOU HAVE JUST HAD YOUR TOOTH/TEETH EXTRACTED WHICH IS SURGERY! BE KIND TO YOURSELF

The initial healing period typically takes from one to two weeks, gum tissue takes about 3-4 weeks to heal and complete healing of the bone can take about 6-8 months depending on your care of this area.

Some discomfort, bleeding and swelling should be expected in the next 24 hours as your mouth heals. Following these simple instructions will normally be all that is needed

As your mouth heals, you can promote faster healing and avoid complications by simply following these instructions.

DO NOT DISTURB THE WOUND: In doing so you may invite irritation, infection and/or bleeding. **The healing process begins** immediately after surgery as your body sends blood to nourish the tooth socket. Complete healing will take one to two weeks and swelling may last 48 hours. Simple pressure from a piece of gauze is usually all that is needed to control the bleeding and to help a blood clot to form in the socket. Sutures may need to be removed in 3-5 days. Keep your fingers and tongue away from this area. No positive or negative pressure like blowing up balloons. Blowing your nose or sneezing violently can dislodge the blood clot and impair healing, so if you have an upper respiratory infection or allergies be sure to take the appropriate medications to treat these conditions.

DO NOT SMOKE OR USE TOBACCO PRODUCTS or ALCOHOL FOR 72 HOURS AFTER THE TOOTH/TEETH Have BEEN EXTRACTED: because smoke can interfere with the healing process, promote bleeding and can cause a very painful situation know as "dry socket". Continuing to smoke during these first few days will slow healing and soft tissue will not be able to begin to fill in the socket to form the blood clot needed for healthy healing. Smoking can also interfere with the bone grow that surrounds the socket resulting in slowing its process of filling in the socket

BRUSHING: For several days after the extraction, it is important to keep the area as clean as possible to prevent infection and promote healing. Do not directly brush the extraction site for the first 3-4 days after surgery to prevent dislodging of the blood clot from the socket. Don't use any toothpaste. Rinsing toothpaste from your mouth could remove the blood clot. Instead this area can be gently and carefully wiped with a clean, wet gauze pad or cloth. After this time you may carefully brush your teeth around this area and floss *gently*.

MOUTHWASH: Avoid all rinsing for 24 hours after extraction. This is to insure the formation of a blood clot. Disturbance of this clot can lead to increased bleeding or the loss of the blood clot. After the first 24 hours you should **VERY GENTLY RINSE** this area 4 times a day using warm salt water (1 tablespoon salt in 8oz of warm water) or mild antiseptic rinses like Breath RX. Rinse very gently after every meal and snack, making sure that the water reaches the extraction site. Rinsing removes food particles and debris from the socket to help promote healing. Chlorhexadine mouth rinse BID for 10 days.

DO NOT SPIT OR SUCK THROUGH A STRAW: This will promote bleeding and may dislodge the blood clot.

BLEEDING: When you leave the office, you will be given verbal instructions regarding the control of

postoperative bleeding. A gauze pad will be placed on the extraction site that you are asked to keep firm pressure on. You will also be asked to change this dressing every 30 to 40 minutes or so depending on the amount of bleeding that is occurring. The gauze should be dampened before placing them over the surgical site. Change gauze only 3-4 times so as not to remove the blood clot. Do not chew on the pack. Do not touch on the extraction site. *It is normal for some blood to ooze from the area of surgery* for 12 hours. We will also give you a package of gauze to take with you to use at home if the bleeding should continue. Should you need to use the gauze at home, remember fold the clean gauze into a pad, thick enough to bite on. Then moisten it and place it directly on the extraction site. Hold it firmly in place, by biting down on the gauze pad or use finger pressure, for about 30-60 minutes. This pressure helps reduce bleeding and permits a clot to form in the tooth socket. If bleeding still continues, moisten a tea bag with water and wrap it in gauze and fold it in half and bite down on it for 30 minutes. Tea contains Tannic Acid which may help to reduce the bleeding. Keep your head elevated with pillows to control bleeding. Use a towel on the pillow the first night,

MEDICATIONS: Some discomfort is normal after surgery. Analgesic tablets i.e. Tylenol or Motrin or non-aspirin can be taken (2) every four hours as needed or as directed by Dr. Peterson. Refer to [pain control](#) article. Prescription medication, which may have been given to you, should also be taken for 2-3 days after surgery or as directed. Don't drive while taking any pain medication due to drowsiness. If antibiotics are prescribed you should carefully follow the instructions and finish the antibiotics until they are completely gone. Avoid aspirin as it may increase bleeding. **A side effect of aspirin is it can increase bleeding by inhibiting blood clotting making it unsuitable following extractions.** To avoid nausea do not take pain medications on an empty stomach.

SWELLING: To prevent swelling, apply an ice pack or a cold towel to the outside of your face in the area of the extraction during the first 12 to 24 hours. Apply alternately, 10-20 minutes on then 10-20 minutes off, for an 6 hour or longer if necessary. Ice is the best restriction to excessive swelling. The more ice, the less swelling. After 24 hours, ice will not stop swelling. Salt water rinses after 24 hours.

SORE JAW JOINT: Your jaw may be sore from holding your mouth open during surgery. This may last 4-5 days. Massage the jaw muscles gently/ Apply moist heat for 10 minutes on/10 minutes off. Eat soft foods. Do not over extend opening your mouth, it can further aggravate this area. There may be some semi-permanent numbing in the area of the extraction. It will decrease in size within the first 6 to 8 weeks and may continue until 6 months after surgery.